

## Proportion of Vaccine-preventable disease cases with complete information, 2012-2015, West Virginia

SURVEILLANCE INDICATOR:		Target	2012	2013	2014	2015
1	The proportion of <b>Haemophilus influenzae invasive disease</b> cases among children under 5 years of age with complete vaccination history.	90%	100% (3/3)	100% (1/1)	67% (2/3)	100% (2/2)
2	The proportion of <b>Haemophilus influenzae</b> isolates from cases under 5 years of age that were serotyped.	90%	100% (3/3)	0% (0/1)	100% (3/3)	100% (2/2)
3	The proportion of <b>measles</b> cases with complete vaccination history.	100%	n/a	n/a	n/a	n/a
4	The proportion of <b>measles</b> cases or chains of transmission that have an imported source.	100%	n/a	n/a	n/a	n/a
5	The proportion of <b>meningococcal</b> cases with complete vaccination history.	90%	75% (3/4)	100% (2/2)	100% (1/1)	100% (2/2)
6	The proportion of <b>meningococcal</b> cases with known serogroup .	90%	75% (3/4)	100% (2/2)	100% (1/1)	50% (1/2)
7	The proportion of <b>mumps</b> cases for which appropriate clinical specimens were obtained and submitted to the laboratory.	80%	0%	67% (2/3)	100% (1)	100% (2/2)
8	The proportion of <b>mumps</b> cases with complete vaccination history.	90%	50% (1/2)	33% (1/3)	100% (1)	100% (2/2)
9	The proportion of <b>pertussis</b> cases from which clinical specimens are obtained.	60%	91% (67/74)	94% (17/18)	100% (18/18)	84% (67/80)
10	The proportion of probable and confirmed <b>pertussis</b> cases meeting the clinical case definition that is laboratory confirmed.	70%	68% (50/74)	89% (16/18)	100% (18/18)	75% (60/80)
11	The proportion of cases confirmed by isolation of <b>B. pertussis</b> by culture.	2%	3% (2/74)	6% (1/18)	11% (2/18)	2% (2/80)
12	The proportion of probable and confirmed <b>pertussis</b> cases with a complete vaccination history.	50%	61% (45/74)	83% (15/18)	94% (17/18)	61% (49/80)
13	The proportion of <b>pneumococcal invasive disease</b> cases among children under 5 years of age with complete vaccination history.	90%	69% (11/16)	88% (14/16)	58% (7/12)	60% (9/15)
14	The proportion of <b>pneumococcal</b> isolates from cases of invasive disease under 5 years of age that are serotyped and tested for antibiotic resistance.	80%	81% (13/16)	69% (11/16)	25% (3/12)	60% (9/15)
15	The proportion of confirmed <b>rubella</b> cases among women of child-bearing age with known pregnancy status.	100%	n/a	n/a	n/a	n/a
16	The proportion of confirmed <b>rubella</b> cases that are laboratory confirmed.	100%	n/a	n/a	n/a	n/a
17	Percentage of <b>varicella</b> cases with complete information for age, vaccination history, and severity of disease.	90%	n/a	n/a	n/a	n/a