

Percentage of Cases with Completed Risk Factor Fields for Selected Food and Waterborne Diseases*, by Region, West Virginia, Cases Reported 2007-2011

<i>Demographic Factors</i> ¹	REGION 1	REGION 2	REGION 3	REGION 4	REGION 5	REGION 6	REGION 7	REGION 8
All Diseases	(N=421)	(N=369)	(N=357)	(N=197)	(N=300)	(N=113)	(N=180)	(N=169)
Sex	100%	99%	99%	100%	99%	99%	99%	100%
Ethnicity	93%	70%	81%	75%	67%	65%	77%	79%
Race	94%	87%	84%	87%	82%	77%	84%	83%
Age	100%	100%	100%	100%	100%	100%	99%	100
Hospitalization	97%	90%	89%	95%	88%	97%	93%	90%
Death	92%	83%	87%	90%	83%	95%	91%	87%

¹ Unknown responses were considered missing

<i>Risk Factors</i> ²	REGION 1	REGION 2	REGION 3	REGION 4	REGION 5	REGION 6	REGION 7	REGION 8
Giardiasis	(N=43)	(N=56)	(N=36)	(N=40)	(N=64)	(N=16)	(N=58)	(N=22)
Hike/Camp/Swim	95%	91%	86%	92%	95%	100%	100%	86%
Drink Untreated Water	95%	91%	86%	92%	97%	94%	100%	86%
Daycare Contact	95%	91%	86%	92%	97%	100%	100%	86%
Travel History	95%	91%	86%	92%	97%	100%	100%	86%
Animal Contact	95%	91%	86%	92%	97%	87%	100%	86%
Source of Water	91%	82%	72%	85%	70%	81%	93%	82%

² Unknown responses were considered a completed answer

* Selected diseases include: Salmonellosis, Campylobacteriosis, Giardiasis, and Shiga toxin-producing *E.coli*