

Percentage of Cases with Completed Risk Factor Fields for Campylobacteriosis by Region,
West Virginia, Cases Reported 2013

	STATE	REGION 1	REGION 2	REGION 3	REGION 4	REGION 5	REGION 6	REGION 7	REGION 8
<i>Demographic Factors</i> ¹	(N=274)	(N=68)	(N=46)	(N=54)	(N=6)	(N=19)	(N=14)	(N=22)	(N=45)
Sex	100%	100%	98%	100%	100%	100%	100%	100%	100%
Ethnicity	89%	97%	91%	80%	83%	89%	100%	77%	84%
Race	91%	99%	91%	80%	100%	95%	86%	82%	98%
Age	100%	100%	100%	100%	100%	100%	100%	100%	100%
Hospitalization	97%	100%	96%	96%	83%	100%	100%	91%	100%
Death	98%	99%	96%	96%	83%	100%	100%	100%	100%
<i>Risk Factors</i> ²									
Eat Raw/Undercooked Poultry	91%	94%	85%	89%	83%	89%	86%	91%	93%
Contact with Poultry	91%	97%	85%	89%	83%	89%	86%	91%	93%
Consume Raw Milk	91%	97%	85%	89%	83%	89%	86%	91%	93%
Consume Raw Dairy Products	91%	97%	85%	89%	83%	89%	86%	91%	93%
High Risk Occupation	86%	91%	78%	81%	83%	95%	86%	86%	96%
Consume Untreated Surface Water	92%	97%	85%	91%	83%	100%	86%	91%	96%
Eat Raw/Undercooked Red Meat	91%	97%	85%	89%	83%	89%	86%	91%	93%
Travel History	92%	97%	85%	91%	83%	100%	86%	91%	96%
Group Activity	90%	99%	80%	89%	83%	95%	86%	91%	87%

¹ Unknown responses were considered missing

² Unknown responses were considered a completed answer, unless case was classified as lost to follow up

Percentage of Cases with Completed Risk Factor Fields for Shiga toxin-producing *E. coli* (STEC)
by Region, West Virginia, Cases Reported 2013

	STATE	REGION 1	REGION 2	REGION 3	REGION 4	REGION 5	REGION 6	REGION 7	REGION 8
<i>Demographic Factors</i> ¹	(N=37)	(N=10)	(N=4)	(N=7)	(N=1)	(N=8)	(N=0)	(N=2)	(N=5)
Sex	97%	100%	100%	86%	100%	100%		100%	100%
Ethnicity	89%	90%	100%	86%	100%	100%		100%	60%
Race	89%	80%	100%	86%	100%	100%		100%	60%
Age	100%	100%	100%		100%	100%		100%	100%
Hospitalization	100%	100%	100%	100%	100%	100%		100%	100%
Death	97%	100%	100%	100%	100%	88%		100%	100%
<i>Risk Factors</i> ²									
Eat Raw/Undercooked Red Meat	81%	90%	100%	71%	100%	88%		100%	100%
Contact with Livestock	86%	90%	75%	71%	100%	88%		100%	100%
Petting Zoo	86%	90%	75%	71%	100%	88%		100%	100%
Occupational Animal Exposure	81%	90%	75%	71%	100%	88%		100%	100%
Eat Fresh Produce	84%	90%	100%	71%	100%	88%		100%	100%
Consume Raw Milk	84%	90%	100%	71%	100%	88%		100%	100%
Consume Raw Dairy Products	86%	90%	75%	71%	100%	88%		100%	100%
Eat Sprouts	86%	90%	75%	71%	100%	88%		100%	100%
High Risk Occupation	70%	70%	100%	43%	100%	88%		0%	80%
Travel History	84%	90%	100%	71%	100%	88%		100%	100%
Group Activity	86%	90%	100%	71%	100%	88%		100%	80%
Consume Untreated Water	86%	90%	75%	71%	100%	88%		100%	100%
<i>Public Health Action</i> ³	(N=33)	(N=9)	(N=3)	(N=5)	(N=1)	(N=8)		(N=2)	(N=5)
Date Education Provided	91%	100%	100%	100%	100%	75%		100%	80%

¹ Unknown responses were considered missing

² Unknown responses were considered a completed answer, unless case was classified as lost to follow up

³ Cases classified as lost to follow up excluded

Percentage of Cases with Completed Risk Factor Fields for Giardiasis by Region,
West Virginia, Cases Reported 2013

	STATE	REGION 1	REGION 2	REGION 3	REGION 4	REGION 5	REGION 6	REGION 7	REGION 8
<i>Demographic Factors</i> ¹	(N=50)	(N=5)	(N=5)	(N=11)	(N=8)	(N=8)	(N=3)	(N=7)	(N=3)
Sex	100%	100%	100%	100%	100%	100%	100%	100%	100%
Ethnicity	84%	100%	80%	73%	75%	100%	100%	71%	100%
Race	86%	100%	80%	73%	100%	100%	100%	86%	100%
Age	100%	100%	100%	100%	100%	100%	100%	100%	100%
Hospitalization	98%	80%	100%	100%	100%	100%	100%	100%	100%
Death	98%	100%	100%	100%	100%	88%	100%	100%	100%
<i>Risk Factors</i> ²									
Hike/Camp/Swim	98%	100%	100%	100%	100%	88%	100%	100%	100%
Consume Untreated Water	98%	100%	100%	100%	100%	88%	100%	100%	100%
Travel History	98%	100%	100%	100%	100%	88%	100%	100%	100%
Recreational Water Exposure	98%	100%	100%	100%	100%	88%	100%	100%	100%
Petting Zoo	98%	100%	100%	100%	75%	88%	100%	100%	100%
Group Activity	90%	100%	100%	100%	88%	75%	100%	86%	67%
High Risk Occupation	90%	100%	60%	100%	100%	75%	100%	100%	67%

¹ Unknown responses were considered missing

² Unknown responses were considered a completed answer, unless case was classified as lost to follow up

Percentage of Cases with Completed Risk Factor Fields for Salmonellosis by Region,
West Virginia, Cases Reported 2013

	STATE	REGION 1	REGION 2	REGION 3	REGION 4	REGION 5	REGION 6	REGION 7	REGION 8
<i>Demographic Factors</i> ¹	(N=194)	(N=46)	(N=33)	(N=37)	(N=12)	(N=23)	(N=16)	(N=14)	(N=13)
Sex	99%	100%	97%	100%	100%	100%	100%	100%	100%
Ethnicity	89%	98%	100%	68%	83%	100%	94%	57%	100%
Race	92%	98%	100%	76%	83%	96%	100%	79%	100%
Age	100%	100%	100%	100%	100%	100%	100%	100%	100%
Hospitalization	97%	100%	100%	92%	83%	100%	100%	93%	100%
Death	95%	98%	97%	92%	83%	96%	100%	93%	100%
<i>Risk Factors</i> ²									
Eat Raw/Undercooked Poultry	86%	93%	76%	78%	83%	96%	88%	93%	85%
Animal Contact (birds, poultry, reptiles)	84%	89%	73%	78%	83%	91%	88%	93%	85%
Eat Raw Eggs	84%	89%	73%	78%	83%	91%	88%	86%	85%
Eat Fresh Shell Eggs	84%	89%	73%	78%	83%	91%	88%	93%	85%
Eat Raw Red Meat	86%	93%	76%	78%	83%	96%	88%	93%	85%
Eat Fresh Produce	83%	89%	73%	78%	83%	87%	88%	86%	85%
High Risk Occupation	73%	70%	67%	73%	67%	74%	69%	100%	77%
Hike/Camp/Swim	86%	96%	76%	78%	83%	96%	88%	86%	85%
Group Activity	84%	91%	73%	78%	83%	91%	81%	93%	77%

¹ Unknown responses were considered missing

² Unknown responses were considered a completed answer, unless case was classified as lost to follow up